

READ THIS LEAFLET

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Smart Flutriafol 500 SC

FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL

Group **3** Fungicide

For control of certain fungal diseases on wheat,
barley and canola when mixed with fertiliser as
per the Directions for Use table



IMPORTANT:
READ THIS LEAFLET BEFORE USE

Crop Smart
better crop protection

Contents: 10, 110 & 1000L

APVMA Approval No.: 65637/51472

Distributed by:
Crop Smart Pty Ltd
ACN 28 093 927 961
2409/4 Daydream Street, Warriewood, NSW 2102
Tel: 1300 783 481 Fax: 1300 783 491

DIRECTIONS FOR USE: ALL STATES

Crop	Disease	Rate per hectare	WHP	Critical Comments																								
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200mL	4 weeks Grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333*</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	333*	70	285	80	250	100	200	120	166												
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																											
60	333*																											
70	285																											
80	250																											
100	200																											
120	166																											
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	100 or 200mL		<p>For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.</p> <table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <th></th> <th>100mL rate</th> <th>200mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>200</td> <td>400 *</td> </tr> <tr> <td>60</td> <td>166</td> <td>333 *</td> </tr> <tr> <td>70</td> <td>143</td> <td>285</td> </tr> <tr> <td>80</td> <td>125 †</td> <td>250</td> </tr> <tr> <td>100</td> <td>100 †</td> <td>200</td> </tr> <tr> <td>120</td> <td>88 †</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)			100mL rate	200mL rate	50	200	400 *	60	166	333 *	70	143	285	80	125 †	250	100	100 †	200	120	88 †	166
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																										
	100mL rate	200mL rate																										
50	200	400 *																										
60	166	333 *																										
70	143	285																										
80	125 †	250																										
100	100 †	200																										
120	88 †	166																										
	Septoria tritici blotch (<i>S. tritici</i>) Take-all (<i>Geaumannomyces graminis</i>)	200mL																										
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	100 or 200mL	<p>* Application volumes over 300mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.</p> <p>† Where application volume is low, product may be diluted with water to improve coverage.</p>																									

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered.

Do not mix seed with treated fertiliser.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Smart Flutriafol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Smart Flutriafol 500 SC Fungicide is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to Smart Flutriafol 500 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Smart Flutriafol 500 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Smart Pty Ltd accepts no liability for any losses that may result from the failure of Smart Flutriafol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection point or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

For REFILLABLE containers (110L, 1000L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and a face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Crop Smart Australia Limited ("Crop Smart") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Crop Smart's skill or judgement in purchasing or using the

same and every person dealing with this product does so at his own risk absolutely. No representative of Crop Smart has any authority to alter these conditions

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

CROP SMART PTY.LTD EMERGENCY CONTACT

1300 783 481

AUSTRALIA WIDE, 24 HOURS