

# Smart Paraquat 250

**DANGEROUS POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
CAN KILL IF SWALLOWED  
DO NOT PUT IN DRINK BOTTLES  
KEEP LOCKED UP

PLACE  
HERE

## HERBICIDE

ACTIVE CONSTITUENT: 250 g/L PARAQUAT  
present as paraquat dichloride

Group **L** Herbicide

For the control of a wide range of grasses and broadleaf weeds as specified in the directions for use table.

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE. THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.



**IMPORTANT:**  
READ THIS LEAFLET BEFORE USING THIS PRODUCT

**Crop Smart**  
better crop protection

APVMA APPROVAL NO: 64588/60109

Contents: 20, 110 & 1000L

Distributed by Crop Smart Pty.Ltd  
ABN: 28 093 927 961  
2409/4 Daydream Street, Warriewood, NSW, 2102  
Tel: 1300 783 481 Fax: 1300 783 491

11/17

## DIRECTIONS FOR USE RESTRAINTS

DO NOT add wetter unless spraying at high volume. Where Smart Paraquat 250 Herbicide is mixed with water at less than 400 mL/100 L of water, add 100 mL Smart Wet 1000 per 100 L of spray.

DO NOT spray plants which are waterlogged, under stress of any kind or covered with soil or dust.

DO NOT spray plants covered with heavy dew, but rain following spraying will not affect results. DO

NOT sow or cultivate for 1 hour after spraying but operations should commence within 7 days.

### For ground application only:

DO NOT use through aircraft, misting machines or hand-held ultra low volume controlled droplet applicators (CDA units).

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Aid to cultivation to minimise cultivation and prepare a clean bed for sowing	Annual grass and broadleaf weed control, Early autumn sowing	Qld, NSW, Vic, SA, Tas, NT, ACT only	1.2 to 1.6 L <sup>1</sup>	Where cultivation follows spraying, it may commence 1 hour after spraying but should be completed within 7 days. Where heavy weed growth is present at spraying a better seed bed will result if cultivation is delayed 3 to 5 days. Use the higher rates for dense, more mature weed stands. Wild Oats must have at least 2 leaves. Where Reglone® is used the lower Smart Paraquat 250 Herbicide rate should be sufficient to control dense mature weeds. <b>Pasture:</b> Remains of old pasture should be reduced by continuous heavy grazing. Remove stock 3 to 5 days before spraying to allow weeds to freshen up.
	Winter, spring and early summer sowing		1.6 to 2.4 L	
	Wild Oats at 2 to 5 leaf stage in autumn/winter	Qld, Vic, SA, Tas, NT only	600 to 800 mL	
		NSW, ACT only	600 mL	
Rice	Annual grass and broadleaf weed control	Qld, NSW, NT only	1.6 L	Pre-sowing.
			800 mL	Post-sowing, pre-crop emergence.

<b>Crop use or Situation</b>	<b>Weeds controlled</b>	<b>State</b>	<b>Rate/ha</b>	<b>Critical Comments</b>
Wild Oat control in Spring Fallows	Wild Oats at 2 to 5 leaf stage	Qld, NSW, NT, ACT only	1.2 to 2 L	Use higher rate for summer growth. Avoid spraying under hot, dry conditions. Best results will be obtained when spraying is carried out in the late evening.
Kikuyu/ Paspalum Pasture	To suppress growth to oversow winter seed	Qld, NSW, ACT only	1.6 or 2.4 L	Use the high rate for February spraying and the low rate in March.
<b>Selective Weed Control</b> Autumn/early Winter Annual Clovers Perennial Clovers	Annual grass and some broadleaf weed control except Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some thistles. For control of these weeds alternative methods such as the spray-graze technique with 2,4-D or MCPA should be considered.	All states	600 mL to 1.2 L  1.2 to 1.6 L <sup>1</sup>	Use the higher rates for dense weed stands.
<b>Late Winter/ early Spring</b> Annual Clovers, Perennial Clovers, Cocksfoot, Perennial Ryegrass, Phalaris, Demeter Fescue only		Qld, NSW, Vic, SA, Tas, NT, ACT only	1.6 to 2.4 L <sup>1</sup>	Use the higher rate in winter/early spring when Barley Grass is present. <b>All applications</b> Graze pastures continuously after the seasonal break to a height of 2 to 4 cm. Remove stock 2 to 3 days before spraying to allow weeds to freshen up. DO NOT apply until clover has reached the 6 leaf stage. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following spring rainfall and growth.

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
<p><b>Late Winter/ early Spring</b> Annual Clovers, Perennial Clovers, Cocksfoot, Perennial Ryegrass, Phalaris, Demeter Fescue only <i>(continued)</i></p>		Qld, NSW, Vic, SA, Tas, NT, ACT only		<p>DO NOT spray clovers which are affected by insect attack, disease or moisture stress and DO NOT use on clover pastures growing in water repellent sands or other situations subject to moisture stress at or immediately following treatment otherwise poor recovery of the clover may result.</p> <p>Use the lower rate for Cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter Fescue. The perennial grasses must be at least 12 months old at spraying.</p> <p><b>DO NOT APPLY TO MEDICS</b></p>
	Yorkshire Fog grass		1.2 L	<p>Apply in early spring to reduce Yorkshire Fog Grass component and increase the clover and desirable grass component. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following spring rainfall and growth. In lower rainfall areas application in mid to late winter may be almost as effective but allow better pasture recovery. If pasture has been grazed allow sufficient time for pasture and Fog Grass recovery before spraying. Apply in spray volumes of 100 to 250 L/ha, the latter for dense or tall, ungrazed pastures. Add Agral at 200 mL/100 L or Smart Wet 1000 at 120 mL/100 L.</p>

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Lucerne Autumn/early Winter  Late Winter/early Spring	Annual grass and some broadleaf weeds	Qld, Vic, SA, WA, Tas, NT only	1.2 to 1.6 L <sup>1</sup>	Use the higher rates for dense weed stands. DO NOT spray lucerne stands under 12 months old. If Mintweed is present use Atrazine 900g/Kg WG Herbicide at 600 g/ha. <b>WARNING</b> - In certain areas, an uncommon species of Barley Grass ( <i>H. glaucum</i> – common Barley Grass is <i>H. leporinum</i> ) resistant to paraquat based products has become established. It may regrow after an initial scorch by Smart Paraquat 250 Herbicide. Where this problem is suspected use Fusilade® for grass weed control. If Smart Paraquat 250 Herbicide has been applied use Fusilade at 1 L/ha after regrowth but before heading.
		NSW only	1.2 L	
		Qld, Vic, SA, WA, Tas, NT only	1.6 to 2.4 L <sup>1</sup>	
		NSW, ACT only	1.2 L <sup>1</sup>	
Perennial grass seed crops Cocksfoot, Perennial Ryegrass, Phalaris, Demeter Fescue only		All states	600 mL to 1.2 L <sup>1</sup>	Use the low rate for Cocksfoot and Perennial Ryegrass and the higher rate for Phalaris and Demeter Fescue. Spray about 4 weeks after a full weed germination following the autumn break. The perennial grasses must be at least 12 months old at spraying.

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Spray topping to reduce seed set. Chickpeas, Faba Beans, Field Peas, Lentils, Lupins, Vetch	Annual ryegrass	NSW, Vic, SA, WA, ACT only	400 or 800 mL	<p><b>As an aid in managing Annual Ryegrass resistance. For use on escapes from a previous herbicide application in the current crop.</b></p> <p>Spray the crop when the Ryegrass is at the optimum stage, that is when the last Ryegrass seed heads at the bottom of the plant have emerged and the majority are at or just past flowering (with anthers present or glumes open) but before haying off is evident - usually October to November.</p> <p>Use of the higher rate in these crops is usually more reliable and gives a greater reduction in seed set.</p> <p>Reduction in crop yield may occur especially if the crop is less advanced relative to the Ryegrass, that is, if crops have a majority of green immature pods.</p> <p>The higher rate may also increase any yield reduction.</p> <p>In practice crop losses in excess of 25% may occur.</p> <p>Apply by ground boom only in 50 to 100 L/ha. Spray with a calibrated boom spray raised to give double overlap at the level of the Ryegrass seed heads.</p> <p>Pressures of 250 to 350 kPa and use of 110015 or 02 nozzles or equivalent will aid coverage.</p>

**DIRECTIONS FOR USE- *continued***

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Spraytopping to reduce seed set. Pastures	Grasses generally (particularly Annual Ryegrass)	All states	400 mL	<p>Heavily graze paddocks during spring flush to encourage even head development. Remove stock 2 to 3 weeks before the anticipated maturity date of the target species.</p> <p>However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed heads at the bottom of the plant have emerged and initial signs of haying off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.</p>
	Barley Grass			<p>Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening of grass seeds, stock (excepting horses) may be returned 24 hours after spraying. Where hardening seeds are present harrow to knock seed from the heads. DO NOT introduce lambs into paddock until safe from risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.</p>
	Saffron Thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Prevention of Annual Ryegrass toxicity	Spraytop - Graze to destroy seed heads	WA only	400 mL	<p>Grazing management as for spray topping above. Remove stock 3 to 4 weeks before the anticipated maturity date. Spray must be applied within 10 days after emergence of the first Ryegrass seed heads. To ensure adequate control of toxin development, <b>heavy continuous grazing is essential from 1 day after spraying</b> until the pasture has completely hayed off. The required stocking rate will vary but must be sufficient to keep all regrowth after spraying completely eaten off to prevent further growth producing new seed heads which could become toxic.</p>
Hay freezing	Maximum retention of protein in standing dry feed	All states	800 mL	<p>Graze paddocks as for spraytopping above. Remove stock 3 to 4 weeks before the anticipated maturity date. Apply prior to commencement of haying off regardless of the grass species involved. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.</p>
General Weed Control-Hops	Annual grasses	Vic, Tas only	1.2 to 1.6 L plus 1.1 kg/ha Atrazine 900 g/Kg granules and/or 750 mL to 1.4 L/ha Reglone <sup>1</sup>	<p>Apply as a directed inter-row spray prior to crop emergence from winter dormancy, using a minimum of 250 L/ha spray volume to ensure good and even coverage of weeds.</p>



**DIRECTIONS FOR USE- *continue***

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Orchards including Bananas, Vineyards	Annual weed control	Qld, Vic, SA, WA, Tas, NT only	1.6 to 3.2 L/sprayed ha + 160 to 320 mL/100 L (a) see below	<p>Spray as necessary for control of annual weeds. Avoid contacting crop foliage. In bananas apply soon after weed emergence and before weeds reach 15 cm in height. Use spraying pressure less than 240 kPa. Avoid chemical contact with roots and peepers near the pseudo stem. Repeat sprays as required.</p> <p>Smart Paraquat 250 Herbicide will not harm trees or vines with mature brown bark if this alone is sprayed. Use the higher rate for dense weed growth.</p> <p>If Fat Hen <i>Chenopodium album</i> or <i>Portulaca</i> spp. are present and Smart Paraquat 250 Herbicide rate is less than the ratio 800 mL/100 L add 200 mL Agral or 120 mL Smart Wet 1000/100 L of spray mix.</p> <p><b>Note:</b> Spot spray rate assumes 1000 L water/ha. For lower water volumes increase dilution rate as below:</p> <p>Water volume 250 L/ha: use 640 to 1280 mL/100 L</p> <p>Water volume 500 L/ha: use 320 to 640 mL/100 L</p> <p>Water volume 750 L/ha: use 210 to 430 mL/100 L</p> <p>OR Measure how much spray is required to cover an area of 100 square metres using your normal application volume. Your dilution rate is 16 to 32 mL of Smart Paraquat 250 Herbicide in this volume.</p>
		NSW only	1.7 L/sprayed ha	

<b>Crop use or Situation</b>	<b>Weeds controlled</b>	<b>State</b>	<b>Rate/ha</b>	<b>Critical Comments</b>
<b>Peanuts</b> Postemergence In crop	Datura spp. (2 to 4 leaf)	Qld, NT only	400 mL	Spray peanuts up to 7 to 8 leaf stage but before majority of plants are flowering. Foliage will be scorched following application but plants recover rapidly. Apply in 200 to 250 L/ha for thorough coverage of weed foliage. A dense canopy of weeds may reduce weed control due to shielding. Add 100 mL Agral or 60 mL Smart Wet 1000/100 L of spray mix. DO NOT spray (on peanuts) under extremely hot dry conditions when peanuts are very small. In environments such as far north Queensland use the lower rates in the range.
	Annual Ground Cherry (2 to 3 leaf), Apple-of-Peru (2 to 4 leaf), Milkweed (2 to 3 leaf)		600 mL	
	Stagger Weed (2 to 3 leaf), Blue Heliotrope (2 to 3 leaf), Wandering Jew (2 to 3 leaf), Anoda Weed (2 to 3 leaf)		800 mL	
	Bellvine (2 to 3 leaf), Common Morning Glory (2 leaf)		1 L	
Potatoes	General weed control (in-crop)	All states	1.2 to 1.6 L <sup>1</sup>	Spray at early crop emergence (no later than 25% emergence of potato shoots). Use the higher rate for dense weed growth.
	Pre-harvest weed control		2.8 L <sup>1</sup>	Spray about 1 week before digging and after tops have died down.
Row Crops, Vegetables, Market Gardens	Pre-planting and pre-crop emergence	All states	1.2 to 1.6 L or 200 mL/100 L <sup>1#</sup>	To control weeds in seed beds. Treat no less than 3 days before sowing or before crop emergence. Use the lower rate for early autumn applications.

**DIRECTIONS FOR USE- *continued***

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Row Crops, Vegetables, Market Gardens <i>(continued)</i>	Post-emergence inter-row weed control	All states	1.2 to 1.6 L or 200 mL/100 L <sup>1#</sup>	Apply after crop seedlings have emerged or when transplanted crops are established. Direct the spray so that it does not touch the crop. Use shielded nozzles or other appropriate means.
	Seedling weeds			Seedling weeds- use the lower rate for early autumn applications.
	Older weeds		2.4 L or 400 mL/100 L <sup>1#</sup>	More mature stages of weed growth.
Sugar Cane (plant and ratoon)	Grasses and some broadleaf weeds	Qld, NSW, NT only	<p><b>Up to 5 cm high</b> 1.2 to 1.6 L per sprayed ha + 1.2 to 1.6 L/ + 275 to 500 g Diuron 900 WG Herbicide</p>	Apply as a broadcast spray over-the-top of plant cane up to the 3 to 4 leaf stage or ratoon cane up to 10 cm high. Cane foliage will be scorched but new leaves will appear in 7 to 10 days.
	Enhancement with Diuron 900 WG Herbicide		<p><b>Up to 10 cm high</b> 1.2 to 1.6 L sprayed ha + 1 kg Diuron 900 WG Herbicide</p>	When spraying cane between 3-4 leaf stage and formation of the true stem: use a directed interspace spray. The Irvin spray boom ( or other similar equipment) is the most suitable to avoid excessive drift onto cane foliage while spraying at the cane bases of plant and ratoon cane. After formation of the true stem, which is resistant to Smart Paraquat 250 Herbicide, the sprayer height can be raised to overlap the spray pattern by raising the dropper to give weed control in the stool. Use the higher rate for dense, more mature weeds. Smart Paraquat 250 Herbicide can be mixed with Atrazine 900 WG Herbicide to give residual weed control when used as a blanket or directed spray- refer to the Atrazine 900 WG Herbicide label for specific rates.# <b>Tank Mixes: See compatibility section</b>

## DIRECTIONS FOR USE- *continue*

Crop use or Situation	Weeds controlled	State	Rate/ha	Critical Comments
Non-Agricultural situations, around sheds, roadways, paths	Annual weed control	All states	1.6 to 4 L/ha or 200 mL/100 L <sup>1#</sup>	Spray to thoroughly wet weed growth. Smart Paraquat 250 Herbicide can be combined with soil residual herbicides Atrazine 900 WG Herbicide to give rapid knockdown and prolonged weed control. Use the higher rate for dense weed growth.
Non-Agricultural situations, around sheds, roadways, paths	Columbus grass	NSW only	<b>Spot Spraying</b> 160 mL/100 L plus 1 L Flupropanate 745 g/L  <b>Boomspray</b> 2.3 to 4.5 L/ha plus 12 to 22 L Flupropanate 745 g/L	-
Firebreak	Knockdown weed growth to eliminate fire hazard or assist firebreak burn	All states	1.6 to 4 L	Apply mid-winter to early summer. Use the higher rate for dense weed growth. After desiccation is complete the sprayed area may be burnt (normally 7 to 10 days after spraying).

<sup>1</sup> if Capeweed or *Erodium* spp. are present: 200 g/L Diquat formulation at 750 mL to 1.5 L/ha (125 mL to 250 mL/100 L for high volume spraying). Use higher rate for plants more than 10 cm diameter.

<sup>#</sup> when applying Smart Paraquat 250 Herbicide at rates less than the ratio 400 mL/100 L add 100 mL Agral or 60 mL Smart Wet 1000/100 L of spray mix. Wetting Agent: (a) Add 170 mL Agral or 100 mL Smart Wet 1000/100 L

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.**

### WITHHOLDING PERIOD

**DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR AT LEAST 1 DAY, OR GRAZE HORSES FOR 7 DAYS AFTER APPLICATION. REMOVE STOCK FROM TREATED AREAS 3 DAYS BEFORE SLAUGHTER.**

**CHICK PEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, VETCH: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

**PLEASE NOTE EXTRA WETTER REQUIREMENTS FOR HIGH VOLUME SPRAYING.**

## **GENERAL INSTRUCTIONS**

This product kills annual grasses and most annual broadleaf weeds (excluding Capeweed) in specified situations and should not be used for any other purpose. Quickly kills green plant tissue on contact. Is immediately inactivated in the soil. At spraying, weeds should be growing vigorously and must not be covered with soil or heavy dew. The principle of selective weed control with this product is that annual weeds are killed but perennial plants and clovers recover after an initial scorch. The control of annual weeds by spraying with this product will allow the desirable perennial species to thicken up at the expense of the weeds. Moisture and fertility should not be limiting at spraying and the proportion of desirable species must be great enough for them to fill in the areas previously occupied by weeds. Long-term weed control can be obtained following the quick knockdown given by this product if it is combined with soil residual chemicals.

1. **DO NOT** use hand-held ultra low volume controlled droplet applicators (CDA units), boomless jets or misting- machines.
2. **Mixing**  
Add the required quantity of product to water in the spray tank and agitate to give even mixing. Agitate again if left standing.
3. **Wetting agent**  
This product contains a wetting agent and additional wetter is not required unless high volume spraying results in excessive dilution of wetter content. This will occur when product rates fall below 400 mL/100 L of spray. Under such circumstances wetter should be added at the rate 100 mL of Agral\* or 60 mL of Smart Wet 1000/100 L of spray mix. Where Fat Hen or Portulaca are present in orchard or vineyard situations, extra wetter should be used when this product ratio is less than 800 mL/100 L. Add wetter at double the above recommendations. Do not use alkaline or anionic wetting agents.
4. **Clean water**  
Mix this product **with clean water only**. Water should be clean and free from clay, silt and algae. Providing it meets this requirement, saline water, water collected from roofs, bore water, dam water and water from creeks may be used.
5. **Application**

### ***(i) Cereals and Broadacre Spraying***

Use a properly calibrated boom spray which should be fitted with flat fan jets and adjusted to a height to give at least double overlap of the spray at the top of the weeds being sprayed. Spraying pressures should be in the range of 200 to 300 kPa. Speed of travel should be in the range of 6 to 15 km/hr. It is essential that a good marking system be used. If a disc marker is used, it must be mounted so as to turn the soil back on to the area sprayed. It is essential to obtain good leaf coverage with the spray and volumes of dilute spray must be adjusted according to density of weed growth. 100 L/ha may be used for seedlings or well grazed weeds up to 2 cm high. For plant height 2 to 5 cm use 150 L/ha and up to 6 to 10 cm use 200 L/ha. Spray volumes may be as low as 50 L/ha (30 L/ha in WA) for weed growth below 5 cm high, or for spray topping and hay freezing.

Equipment must be appropriate to this volume, properly calibrated and fitted with spraying tips designed to give droplets in the 200 to 250  $\mu$  Volume Median Diameter range.

### ***(ii) High Volume Application***

Higher volumes will generally be required to give good coverage of weed growth in situations other than those specified under cereals and other broadacre crops.

(iii) Wash spray equipment with clean water immediately after use. This product is highly corrosive to metals, particularly galvanised iron and aluminium and should not be left for long periods in tanks or equipment made of these materials.

## 6. Compatibility

This product combines satisfactorily with the soil active herbicides Atrazine 900 WG Herbicide, Diuron 900 WG Herbicide and Simazine 900 WG Herbicide where prolonged weed control is required as well as a quick knockdown. This product is compatible with Smart Wet 1000 wetting agent, Agral, BS1000, Reglone®, Spray Seed® 250, Banvel® 200, Banvel® M, MCPA Amine (no more than 1 L/800 mL Smart Paraquat 250 Herbicide), CHLORSULFURON 750 WG Herbicide, Oryzalin/Trifluralin 250 EC Herbicide, Smart Tri-allate 500 EC Herbicide, and Smart Trifluralin 480 EC Herbicide.

**TANK MIXTURES:** Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products.

## 7. Spraying Conditions

Avoid spraying plants under stress from waterlogging, frost, drought etc. or covered with dust and soil. Results will be better if application is made in dull weather or at the end of the day. Light rain following spraying will not affect results. Avoid drift into neighboring crops.

## Resistant Weeds Warning

<b>GROUP</b>	<b>L</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Smart Paraquat 250 Herbicide is a member of the bipyridyls group of herbicides. Smart Paraquat 250 Herbicide has the inhibitor of photosynthesis at photosystem I mode of action. For weed resistance management Smart Paraquat 250 Herbicide is a Group L herbicide. Some naturally occurring weed biotypes resistant to Smart Paraquat 250 Herbicide and other Group L herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by Smart Paraquat 250 Herbicide or other Group L herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Crop Smart Pty Ltd accepts no responsibility for any losses that may result from failure of Smart Paraquat 250 Herbicide to control resistant weeds.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto susceptible plants/crops, cropping lands or pastures. This formulation should not be applied on or near water which is used for irrigation purposes.

### PROTECTION OF LIVESTOCK

Domestic pets and poultry - keep away from treated areas. This formulation should not be applied on or near water which is used for livestock watering.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

This formulation should not be applied on or near water which is used for human consumption, livestock watering or irrigation purposes or water used for commercial or recreational fishing.

### STORAGE AND DISPOSAL

#### (20L only)

Store in the closed original container, in a dry, cool well-ventilated area out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

Empty containers should not be burnt. DO NOT burn empty containers or product.

### **Refillable containers (1000L only)**

Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **Envirodrum Mini Bulk Container (110L and 200L)**

Store in the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of Crop Smart Pty Ltd.

### **SAFETY DIRECTIONS**

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks eyes, protect eyes while using. Avoid contact with eyes, skin and clothing. When opening the container and preparing Smart Paraquat 250 Herbicide for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product, remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist.

DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

### **SPRAY APPLICATION**

DO NOT work in spray mist. DO NOT continue to use if skin irritation or nose bleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice. When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended, but in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer. Avoid contacting vegetation wet with spray, but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

### **FIRST AID**

If poisoning occurs get to a doctor or hospital quickly. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Smart Paraquat 250 Herbicide is available from Crop Smart Pty Ltd on request. Call Customer Service on 1300 783 481.

## CONDITIONS OF SALE

Crop Smart Pty Ltd ('Crop Smart') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Crop Smart's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Crop Smart has any authority to add to or alter these conditions.

**UN 3016**  
**BIPYRIDILIUM,**  
**PESTICIDE, LIQUID, TOXIC (CONTAINS PARAQUAT)**  
**HAZCHEM CODE: 2X**  
**PACKAGING GROUP: III**

<p>IN A TRANSPORT EMERGENCY <b>DIAL 000</b> POLICE OR FIRE BRIGADE</p>
<p><b>CROP SMART PTY.LTD EMERGENCY CONTACT</b> <b>1300 783 481</b> <b>AUSTRALIA WIDE, 24 HOURS</b></p>

