

Smart

Trifluralin 480 EC

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PEEL
HERE

HERBICIDE

ACTIVE CONSTITUENT: 480 g/L TRIFLURALIN
SOLVENT: 480 g/L LIQUID HYDROCARBON

Group **D** Herbicide

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.



**IMPORTANT: READ THE ATTACHED
LEAFLET BEFORE USING THIS PRODUCT**

Crop Smart
better crop protection

APVMA Approval Number: 61325/B/1007

Contents: 20, 110 & 1000L

Distributed by
Crop Smart Pty Ltd
ACN: 093 927 961
2409/4 Daydream Street
Warriewood, NSW 2102

STORAGE AND DISPOSAL

(20L and 200L only)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

Envirodrums: 110L Mini Bulk Returnable Containers

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Crop Smart Pty.Ltd.

Refillable Containers: 1000L

Store in the closed, original container in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

Crop Smart Pty Ltd ('Crop Smart') will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Smart is limited to the replacement of the goods or (at the option of Crop Smart) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Smart within thirty days of delivery.

DIRECTIONS FOR USE

1. FIELD CROPS

Situation & Crop	Weeds	State	Rate / Soil Type			Critical Comments
			Light	Medium	Heavy	
Chickpeas	Annual Ryegrass, Paradoxa Grass (Canary Grass), Wireweed (Hogweed), Black Pigweed. Suppression of Climbing Buckwheat (Black Bindweed), soil surface Wild Oats	Qld only	1.25 - 1.7L/ha	1.25 - 1.7L/ha	1.25 - 1.7L/ha	Use 1.25L/ha when applying immediately prior to sowing. Use 1.7L/ha when applying to dry soil before the planting rain.
	Annual Ryegrass, Wireweed (Hogweed), Deadnettlles, Wild Oats	Vic only	800mL/ha plus 1.6L/ha Smart Tri-Allate 500 EC Selective Herbicide			Incorporate as per recommendations for wheat, barley and tritiale.
	Red & White Fumitory, Rough Poppy, Wireweed, Annual Ryegrass, Barley Grass, Canary Grass, Sand Fescue, Suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr weed, Brome Grass, Cereal Oats and soil surface Wild Oats	SA only	1.25L/ha	1.25L/ha	1.25L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Incorporate as per Incorporation Table 6.
	Amaranthus,	NSW,	1.2 –	1.5L/ha	1.7L/ha	Apply from 4 weeks up

	Annual Ryegrass, Barnyard Grass, Caltrop, Crab Grass, Paradoxa Grass (Canary Grass), Pigweed, soil surface Wild Oats, Winter Grass, Wireweed, Suppression of Fumitory	ACT only NSW, ACT, Qld only	1.5L/ha			to just prior to sowing. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Adzuki Beans, Cowpeas, Lablab, Mung Beans, Borlotti Beans, Red Kidney Beans						
Faba Beans	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivy Leaf Speedwell, Mustards, Turnips, Wireweed. Suppression of Brome Grass, Soursob and soil surface Wild Oats	SA, WA only	800mL/ha plus 1 – 1.5L/ha Simazine Flowable (500g/L)			Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For full reliable results, significant rainfall (20 or 30mm) is necessary within 2-3 weeks of application.
Pigeon Peas	Amaranthus, Barnyard Grass, Canary Grass, Crowsfoot Grass, Pigweed, Spiny Burrgrass, Summer Grass, soil surface Wild Oats,	NSW, ACT only	1.2L/ha	1.5L/ha	1.7L/ha	Apply between 4 weeks and just before sowing. Refer to Incorporation Table 3, 4 or 6 for suitable method of incorporation.

	Wireweed (Hogweed), Suppression of Yellow Vine (Caltrop), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass					
Lentils	Annual Phalaris, Annual Ryegrass, Wild Oats, Wireweed	NSW, ACT only	800mL/ha	1.2L/ha	1.2L/ha	Apply 1 to 4 weeks before sowing.
	Fumitory – Red and White, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue	SA only	1.25L/ha	1.25L/ha	1.25L/ha	Apply 1 to 4 weeks before sowing.
Navy Beans	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop (Bullhead, Yellow Vine), Crab Grass, Mossman River Grass (Innocent Grass), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather),	All States	1.2L/ha	1.5L/ha	1.7L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Soybeans			1.2L/ha	1.7L/ha	2.3L/ha	

	Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)					
Vetch	Annual Ryegrass, Deadnettle, Wireweed, soil surface Wild Oats, suppression of Brome Grass, Rough Poppy, Speedwell, Three Cornered Jack, Yellow Burr Weed, Sheepweed	SA, WA only	1.7L/ha	1.7L/ha	1.7L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Refer to Incorporation Table 6 for method of incorporation.
Cotton	Annual Ryegrass, Barnyard Grass, Canary Grass,	Qld, ACT, NSW, WA only	1.2L/ha	1.7L/ha	2.3L/ha	Spray between 6 weeks and just before sowing takes place. Refer Incorporation Table 1 & 2 for method of incorporation.
Legume Seed Crop Establishment - Annual Medics - Clover (Berseem, Red, Strawberry, Sub and White)	Caltrop (Bullhead/ Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus),	NSW, ACT, SA, WA, Vic, Tas only	1.2L/ha	1.2L/ha	1.7L/ha	Autumn Sowing – Apply from 4 weeks to 7 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation.
- Lucerne for hay and	Redshank (Prince of	All States	1.7L/ha	1.7L/ha	1.7L/ha	Spring Sowing – Apply between 4 weeks and 3 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation.

seed crop	Wales Feather), Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed), Black Pigweed (Qld only), From					In both cases seedling disease, cold weather, excessive moisture, high salt concentrations and drought could weaken crop seedlings and damage could occur from the use of this product. Temporary crop suppression could result.
Linseed	seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed	NSW, ACT, SA, Wa, Vic only	1.2L/ha	1.5L/ha	1.7L/ha	Spray 2-4 weeks before sowing. Sowing depth should be 1.3 to 2.5cm. Deeper sowing may result in some stand reduction. Refer Incorporation Table 6 for method of incorporation.
Peanuts	Grass (Urochloa)	WA, Qld only				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Peas		All States				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 or 11 for method of incorporation.
Canola, Safflower						
Sugar cane - Early Season - Late Season		Qld, NSW only	3.0L/ha 2.3L/ha	3.0L/ha 2.3L/ha	3.0L/ha 2.3L/ha	Apply to plant cane after emergence to "out of hand" stage. Apply to ratoon cane immediately after harvest. Refer Incorporation Table 3, 7 or 10 for suitable method of incorporation.

1.FIELD CROPS continued

Situation & Crop	Weeds	State	Rate / Soil Type			Critical Comments
			Light	Medium	Heavy	
Sunflowers	Refer to weeds on previous page	All States	1.2L/ha	1.5L/ha	1.7L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for method of incorporation.
Lupins			Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 for method of incorporation.			
	Annual Grasses and Broadleaf Weeds	NSW, ACT, Vic, Qld only	800mL/ha plus 3L/ha of Simazine Flowable (500g/L)			Use a low volume boom applying 50-100 Litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing the crop. Incorporate within 4 hours of application. Do not apply to a ridged soil.
	Capeweed, Turnip, Radish, Doublegee and Suppression of Annual Ryegrass and soil surface Wild Oats	WA only	1.25L/ha plus 1 – 1.5L/ha of Simazine Flowable (500g/L)			Rate for Yellow Sands.
	As above plus suppression of Brome Grass		1.25L/ha plus 2L/ha of Simazine Flowable (500g/L)			Rate for all other soil types. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For Simazine to be effective sufficient rainfall (20-30mm) to wet the soil through

						the weed root zone is necessary within 2-3 weeks of application. Results with Simazine can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into dry or low moisture seed beds.
	Capeweed, Doublegee, Wild Radish, Wild Turnip plus suppression of Annual Ryegrass, soil surface Wild Oats and Brome Grass		1.25L/ha plus 2L/ha of Diuron Flowable (500g/L)			DO NOT use on white or grey sands as severe crop damage may result. Use tank mix of Diuron & Trifluralin where Annual Ryegrass is present. Apply pre-sowing Stage.
	Red & White Fumitory, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue, suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats, soil surface Wild Oats	SA only	1.25 – 1.7L/ha	1.25 – 1.7L/ha	1.25 – 1.7L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 for method of incorporation.
	Above weeds plus Capeweed, Common Fumitory, Geranium, Indian Hedge Mustard, Sheepweed, Shepherd's Purse,	SA only	1.25L/ha to 1.7L/ha plus 2 to 4L/ha of Simazine Flowable (500g/L)			Use a low volume boom applying 50-100 Litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing the crop. Incorporate within 4 hours of application. Do not apply to a

	Toadrush, Turnips, suppression of Ice Plant, Soursob					ridged soil.	
Tobacco	Summer Grass, Crowsfoot Grass, Red Natal Grass, Love Grass, Button Grass, Rhodes Grass, Pigweed	Qld only	800mL/ha	1.2L/ha	-	Apply to soil 3-4 weeks prior to transplanting. The longer period to be used for applications made during June & July. Incorporate to a depth of 10cm.	
	Crowsfoot Grass	NSW, ACT only	800mL/ha	1.2L/ha	1.2L/ha	Apply to light sandy soil 14 to 21 days before transplanting. Do not incorporate to a depth greater than 6cm. Apply to loam (medium soil) 14 to 21 days before transplanting. DO NOT incorporate to a depth of greater than 6cm.	
Wheat, Barley & Triticale	Annual Ryegrass, Wireweed (Hogweed), Phalaris spp.	NSW, ACT, Vic only	800mL/ha	800mL/ha	800mL/ha	Apply 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Incorporation Table 6 for method of incorporation.	
	1. Pre- Sowing Only	Fumitory					WA only
	2. Pre- Sowing & Post Sowing (self- mulching soils)	Canary Grass					Vic only
		As above, except for Fumitory					Vic only
						Pre-sowing – Apply more than 4 weeks before sowing to prevent crop damage. Post Sowing – Apply within 2 days after sowing to well prepared seedbed. Refer Incorporation Table 6 for suitable	

						method of incorporation.
Wheat	Annual Ryegrass, Paradoxa Grass (Canary Grass), soil surface Wild Oats, Wireweed (Hogweed)	Qld only	800mL/ha	800mL/ha	800mL/ha	On non self-mulching soils apply 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. On self-mulching soils, as above except apply more than 4 weeks before sowing to prevent crop damage. Refer Incorporation Table 6 for method of incorporation.
Barley						Apply to self-mulching and non self mulching soils from 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. Refer Incorporation Table 6 for suitable method of incorporation.
Wheat, Triticale, Rye	Annual Ryegrass, Red & White Fumitory, Phalaris spp., Wireweed, suppression of Deadnettle, Rough Poppy, Yellow Burr Weed	SA only				Apply 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Incorporation Table 6 for method of incorporation.
Barley	As above for SA plus Sand Fescue and suppression of Brome Grass		1.25L/ha	1.25L/ha	1.25L/ha	
Wheat & Triticale only	Annual Phalaris	NSW, ACT only	800mL/ha plus 20g/ha of Chlorsulfuron 750 WG Herbicide			If possible, spray and incorporate into the soil in one operation. If this is not possible

				incorporation should take place within 4 hours of spraying. Delay may cause inferior weed control.
--	--	--	--	--

2. FIELD CROPS: FOR USE IN NO-TILL/MIN-TILL CROPPING, PRE-SOWING OR INCORPORATED BY SOWING (IBS). (Using Incorporation Table 13)

Situation & Crop	Weeds	State	Rate	Critical Comments
Wheat, barley and triticale	Annual Ryegrass, Wireweed, Phalaris spp, Fumitory	WA, SA, Vic, NSW, ACT only	1.5L-2L/ha	Use Higher rate on Lighter Sandy and Sandy Loam Soils. DO NOT use on heavy soils. Use with Knife/Blade point sowing equipment. Use higher rate for heavier stubble coverage. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Refer Incorporation Table 13 for method of incorporation. Application can occur 0-24 hrs prior to incorporation by sowing. For best results apply as close as possible to sowing (within 12 hours). Application 12-24 hours before sowing may be more adversely affected by above average soil moisture, warm winter temperatures and high weed seed densities. These factors individually or combined may reduce final weed control levels.
Chickpeas		WA Only	1.25-1.7L/ha plus 1.1kg/ha Simazine WG Herbicide (900g/Kg)	Incorporate as per Incorporation Table 13.

3. VEGETABLES, ORCHARDS AND VINEYARDS

Situation & Crop	Weeds	State	Rate / Soil Type			Critical Comments
			Light	Medium	Heavy	
Transplants Only Broccoli, Cabbage, Cauliflowers, Tomatoes	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop (Bullhead Yellow Vine), Crab Grass, Mossman River Grass (Innocent Grass),	All States	1.2L/ha	1.7L/ha	2.3L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Direct Seeded Only Broccoli Brussels Sprouts,						

Cabbage	Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)					Apply to new planting during pre plant cultivation. Apply to established crops in spring after weeds and green manure crop has been ploughed into ground. Refer Incorporation Table 8 or 9 for suitable method of incorporation.
Cauliflower		Vic only				
Carrots		All States				
Chicory		Vic only				
Green Beans		All States	1.2L/ha	1.5L/ha	1.7L/ha	
Orchards and Vineyards		Qld, SA, WA, Vic, Tas only	1.2L/ha	1.7L/ha	2.3L/ha	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

THIS PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 4 HOURS OF APPLICATION EXCEPT WHERE THE CROP IS SOWN WITH MINIMUM TILLAGE SOWING EQUIPMENT (FITTED WITH KNIFE POINTS OR BLADES LESS THAN 12MM WIDE, USUALLY WITH PRESS WHEELS) WHERE APPLICATION MAY OCCUR UP TO 24 HOURS BEFORE INCORPORATION BY THE SOWING PROCESS.

INCORPORATION TABLE

1. Prior to furrowing out: - 2 workings at an angle required using Offset or Tandem disc harrows.
2. After furrowing out: - 2 workings required using GoDevil discs or Lilliston cultivators set at 10 cm depth.
3. Rotary Hoe: - 1 working required at 5-7.5cm depth. Sugar Cane: - 7.5-13cm depth.
4. Offset or Tandem Disc Harrows:- (preferably with spiked harrows in tandem) 2 workings at an angle required at 7.5-15cm depth at 6.5-10km per hour.
5. Heavy Diamond or Stump Jump Harrows: - (weighted 20-30kg per section) at 10-13 km/h speed. Then cross work with offset or tandem disc harrows set to 7.5-15cm depth at speed 6.5-10km per hour.
6. Weighted Heavy Diamond or Stump Jump Harrows: (weighted with 20-30kg per section) at 10-13km/hr. Cross work with combine at 5-7.5cm depth at speed of 10-13km/hr.
7. Disc Ratoon Cultivator: - 2 workings needed with discs and cultivator set at 7.5-13cm depth.
8. Offset or Tandem Disc Harrows:- set at 7.5-15cm depth. A second discing is required working in opposite direction with discs set to throw treated soil into tree or vine row.
9. Rotary Hoe: - 1 working needed at 5-10cm depth.
10. Offset Discs (Bumpers): - 2 workings needed at depth of 7.5-13cm.
11. Incorporation by sowing (IBS) on suitably prepared seedbed with heavy diamond harrows trailing or as a separate operation.

12. Post-sowing/Pre-emergence: Use heavy diamond harrows cross working at right angles to the direction of sowing. Do not attempt this method of incorporation on poorly prepared, clumpy or cloddy soils.
13. Incorporation by Sowing (IBS) with knife or blade points. Use press wheels to avoid dragging treated soil back into the seed furrow. Maintain slow to moderate speed to ensure that soil throw is not into adjacent furrows." Note a. - Knife or blade point systems can result in poor weed control in the seed furrow as chemical displacement from this zone occurs. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. b. A Knife or blade point is 12mm or less, has no wings, inverted T or blade, and is generally placed on a minimum 20cm tyne spacing.

MIXING

This product is an emulsifiable concentrate which mixes readily with water. Add the recommended amount to the spray tank during filling operation and apply 70-450L of water/ha (broadcast basis) dependent on soil type and stubble coverage level. For minimum tillage/stubble retention seeding systems use of higher water volumes may help reduce the impact of stubble.

Ensure adequate agitation is continued throughout the operation. Leaving the made up spray mixture for long periods of time without agitation is not recommended.

Under hot conditions or where possible spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Use properly calibrated standard low pressure (170-340 kilopascal) boom type sprayer with fan tips.

CONDITIONS FOR BEST RESULTS

This product must be thoroughly incorporated as recommended. Soil should be well worked and free of weeds at time of application. Product effectiveness may be reduced by inadequate incorporation, high organic matter, excess clods, crop or trash residues, stones or other foreign matter and in areas of unnaturally high weed seed population such as header tracks or livestock rest areas.

Trifluralin is volatile and disappears from exposed surfaces. Loss is hastened by high temperatures, winds or warm moist soil.

Integrated Weed Management:

The use of Integrated Weed Management techniques in conjunction with Trifluralin 480 are always recommended. Agronomic practices that reduce the weed seed bank in the soil prior to the use of Trifluralin 480 will result in higher weed control levels from Trifluralin 480. Failure to use Agronomic and Integrated Weed Management practices that reduce the weed seed bank in the soil will result in higher weed seed populations. Paddocks with excessively high weed seed banks may have sufficient weed numbers surviving such that final weed control may be considered below a commercially acceptable level and additional herbicide treatments may be necessary. The use of Integrated Weed Management techniques will also reduce the potential for the development or survival of Group D herbicide resistance weed biotypes.

WILD OATS

Germinating wild oat seeds lying on soil surface will be controlled. Therefore, specific wild oat control is only possible with shallow cultivation. Poor control will occur on self mulching soils and all soil types where deep cultivation is practised.

COMPATIBILITY

This product may be mixed in the spray tank with:

Herbicides: Smart Tri-Allate 500 EC Selective Herbicide, Chlorsulfuron 750 WG Herbicide, Smart Triasulfuron 750 WG Herbicide, Diuron Flowable (500g/L), Simazine Flowable (500g/L), Smart Paraquat 250 Herbicide, Spray*seed*, Broadstrike*, Spinnaker*, Bladex*, Sencor*/Lexone*, Smart Glyphosate 450 SL Herbicide

Insecticides: Chlorpyrifos 500EC, Endosan EC.

NOTE:

1. Information on compatibility is understood to be correct at the time of publication, however products may vary from time to time, therefore a small scale compatibility test should be carried out before mixing in the spray tank.
2. Observe any mixing sequence instructions for tank mix products.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise, spray tanks, pumps, lines and nozzles should be thoroughly rinsed several times with clean water following application. Spraymate Tank & Equipment Cleaner is suitable for this purpose and will also remove Trifluralin stains.

RESISTANT WEEDS WARNING

GROUP	D	HERBICIDE
--------------	----------	------------------

Smart Trifluralin 480 EC Herbicide (Trifluralin 480) is a member of the DINITROANILINES group of herbicides. Trifluralin 480 has the inhibitors of tubulin formation mode of action. For weed resistance management Trifluralin 480 is a Group D herbicide. Some naturally-occurring weed biotypes resistant to Trifluralin 480 and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Trifluralin 480 or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Crop Smart Pty Ltd accepts no liability for any losses that may result from the failure of Trifluralin 480 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT use in high winds. DO NOT exceed rates specified, to avoid crop damage. DO NOT plant sensitive grasses such as oats, sorghum, millets, phalaris spp., ryegrass or wheat for 12 months following the use of this product except where wheat follows wheat or other winter crops. DO NOT plant oilseed poppies when a detectable residue of Trifluralin is present in the soil. Levels as low as 0.02ppm may interact with other unfavourable factors (moisture, stress, disease etc.) to reduce poppy growth and vigour. DO NOT apply to orchards and vineyards after first flush of growth or when residues can lodge on or in fruit. Reduced germination of wheat and barley may occur due to combination of following circumstances and the use of this product:

- Short coleoptile cultivars
- Use of seed dressings (except Vitavax*)
- Shallow or uneven seedling depth.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry well-ventilated area, out of direct sunlight. Do not store below 5°C. Extended storage below 5°C can result in the formation of crystals on the bottom of the container. If crystallisation does occur, store the container on its side at room temperature and rock occasionally until crystals re-dissolve. Ensure any crystals are dissolved before adding to the spray tank.

20L and 200L containers only:

Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty

packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

Refillable containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container (110L only)

Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Crop Smart Pty Ltd.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Telephone Australia: 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

Crop Smart Pty Ltd ('Crop Smart') will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Smart is limited to the replacement of the goods or (at the option of Crop Smart) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Smart within thirty days of delivery.

** Not a registered trademark of Crop Smart Pty Ltd*



**IN A TRANSPORT EMERGENCY DIAL 000
POLICE OR FIRE BRIGADE**

**CROP SMART PTY LTD EMERGENCY CONTACT
1300 783 481 AUSTRALIA WIDE, 24 HOURS**