

# DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Smart

# Scavenge Sterilised Grain Mouse Bait

ACTIVE CONSTITUENT : 25 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT**

Crop Smart Pty. Ltd  
ABN 28 093 927 961  
2409/4 Daydream Street, Warriewood NSW 2102  
Tel: 1300 783 481 Fax: 1300 783 491



Batch Number:

Date of Manufacture:

Scan QR Code to  
access SDS  
on mobile device



APVMA Approval Number: 90334/130511

**Crop Smart**  
better crop protection

# DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Smart

# Scavenge Sterilised Grain Mouse Bait

ACTIVE CONSTITUENT : 25 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

**CONTENTS: 125kg - 500kg**

Crop Smart Pty. Ltd  
ABN 28 093 927 961  
2409/4 Daydream Street, Warriewood NSW 2102  
Tel: 1300 783 481 Fax: 1300 783 491

APVMA Approval Number: 90334/130511

**Crop Smart**  
better crop protection



## **DIRECTIONS FOR USE:**

### **RESTRAINTS**

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent

DO NOT apply to the outer 50m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

Situation	Pest	Rate	Critical Comments
<p>Grain crops, legume crops, canola, safflower, nut crops, and pasture.            (Note: for all situations - only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).</p>	<p>Mice</p>	<p>1 kg/ha to achieve an even coverage of 2-3 grains/m<sup>2</sup></p>	<p>Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m<sup>2</sup>. It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals.</p> <p>Smart Scavenge Sterilised Grain Mouse Bait will provide a high level of control in most situations. However, in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment after 14 days may therefore be required in certain circumstances.</p> <p>It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control.</p> <p>Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.</p>

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

**WITHHOLDING PERIOD**

DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK**

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock.

It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

DO NOT contaminate dams, streams, rivers or watercourses with the product or used containers.

**STORAGE AND DISPOSAL**

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Shake and empty contents into application equipment. Do not dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**Metal drum** - If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

**SAFETY DIRECTIONS**

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. DO NOT inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits, wear elbow-length PVC gloves and a full-face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26; New Zealand 0800 764 766). Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxyviva or one-way mask. Resuscitate in a well-ventilated area.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet (SDS), which is available from the supplier.

**NOTICE TO BUYER**

Crop Smart Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Crop Smart's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk.



**Additional GHS Statement**

Contact with acid liberates very toxic gas.  
In contact with water releases flammable gases.  
Toxic if swallowed.  
Very toxic to aquatic life with long lasting effects.

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

**EMERGENCY CONTACT CHEMWATCH  
1800 951 288  
AUSTRALIA WIDE, 24 HOURS**

